

## Pocket Gophers

In some areas, the name gopher is applied to a variety of mammal species including ground squirrels and moles. True pocket gophers are burrowing rodents that get their name from their fur-lined cheek pouches, or pockets. These pockets are used, like a squirrel's, for carrying food. However, the pockets on a gopher open on the outside and turn inside out for emptying and cleaning.

Pocket gophers are well-equipped for a digging, tunneling lifestyle, with large-clawed front paws, small eyes and ears, and sensitive whiskers that assist with movement in the dark (Fig. 1). Their pliable fur and sparsely haired tails—which also serve as a sensory mechanism—help gophers run backward almost as fast as they can run forward. Their large front teeth are used to loosen soil and rocks while digging, as well as to cut roots (Fig. 2).

The pocket gopher's short fur is a rich brown or yellowish brown, but also may be grayish or closely resemble the local soil color.

Two species of pocket gophers occur in Washington: the **Northern pocket gopher** (*Thomomys talpoides*) is the smallest and most widespread, occupying much of eastern Washington. Adults of this species measure 8 inches in length, including their 2-inch tail. A subspecies, called the **Brush Prairie pocket gopher**, occurs in Clark County (see "Legal Status"). The **Mazama (Western) pocket gopher** (*Thomomys mazama*) is the only pocket gopher in most of western Washington—in the Olympic Peninsula and the southern Puget Sound area. Adults measure 8 inches in length, including their 2½-inch tail (see "Mazama Pocket Gopher Conservation").

Pocket gophers can be a problem for homeowners, but they actually benefit the soil and vegetation in many areas. Unfortunately, the positive effects are not as visible as the mounds pocket gophers create in lawns and pastures.



**Figure 1.** Pocket gophers (Northern pocket gopher shown here) are stout-bodied rodents with small ears and eyes and large clawed front paws. Their large front teeth are used to loosen soil and rocks while digging, as well as to cut and eat roots.

(From Christensen and Larrison, *Mammals of the Pacific Northwest: A Pictorial Introduction*.)


## Facts about Washington's Pocket Gophers

### Food and Feeding Habits

- Unlike moles, which mostly eat insects and other invertebrates, pocket gophers only eat vegetation.
- Gophers eat roots, bulbs, and other fleshy portions of plants they encounter while digging underground.
- Gophers also eat the leaves and stems of plants around their tunnel entrances and can pull entire plants into their tunnels.
- In areas with a snowpack, gophers will gnaw on bark several feet up a tree or shrub.
- Because gophers obtain sufficient moisture from their food, they don't need a source of open water.

## Reproduction and Social Structure

- Pocket gophers breed from early spring to early summer, resulting in one litter of three to seven young per year.
- The nest chamber is located in the pocket gopher's burrow system, is about 10 inches in diameter, and is lined with dried vegetation
- The young develop quickly, remain in the nest for five to six weeks, and then wander off above ground to form their own territories.



- Pocket gophers are solitary except during the breeding season or when females have young with them.
- Densities of northern pocket gophers have been found to range from 2 to 20 gophers per acre, depending on food availability, species, and ages of the gophers.

## Mortality and Longevity

- Coyotes, domestic dogs and cats, foxes, and bobcats capture gophers at their burrow entrances; badgers, long-tailed weasels, skunks, rattlesnakes, and gopher snakes corner gophers in their burrows. Owls and hawks capture gophers above ground.
- A deep snowpack can result in high gopher mortality. If the snow melts rapidly it saturates the ground and floods the burrows.
- Pocket gophers live one to two years and the majority of the population consists of young adults.

## The Benefits of Pocket Gophers

A typical pocket gopher can move approximately a ton of soil to the surface each year. This enormous achievement reflects the gopher's important ecological function.

Their tunnels are built and extended, then gradually fill up with soil as they are abandoned. The old nests, toilets, and partially filled pantries are buried well below the surface where the buried vegetation and droppings become deep fertilization. The soil thus becomes mellow and porous after being penetrated with burrows. Soil that has been compacted by trampling, grazing, and machinery is particularly benefited by the tunneling process.

In mountainous areas, snowmelt and rainfall are temporarily held in gopher burrows instead of running over the surface, where they are likely to cause soil erosion.

Surface mounds created by gophers also bury vegetation deeper and deeper, increasing soil quality over time. In addition, fresh soil in the mounds provides a fresh seedbed for new plants, which may help to increase the variety of plants on a site.

Many mammals, large birds, and snakes eat gophers and depend on their activities to create suitable living conditions. Salamanders, toads, and other creatures seeking cool, moist conditions take refuge in unoccupied gopher burrows. Lizards use abandoned gopher burrows for quick escape cover.

## Viewing Pocket Gophers

Although pocket gophers are active year-round and at all hours of the day, their underground lifestyle makes them difficult to observe.

If you are patient, you may be able to watch a pocket gopher feed above ground, or see their food being taken underground. The Mazama pocket gopher spends some time above ground, generally at night and on overcast days.

When sitting in a grassy area, keep your eyes and ears alert for the sight and sound of a wiggling clump of grass, wild flowers, or similar vegetation. You might see the entire plant slowly disappear below ground. A few minutes later, the same gopher may venture a body length's distance from its tunnel opening to alertly feed or gather food.

When gathering food above ground, a pocket gopher will cut vegetation quickly, cram as much as possible into its external pouches (or pockets), and then disappear below ground. It may reappear in a few minutes, gather more food, and disappear to consume the food underground or store it away for later.

Pocket gophers live in extensive burrow systems, which they use for locating food, rearing young, storing food and droppings, and escaping predators. Burrow systems are a closely regulated microenvironment, and gophers will plug any openings in the system within 24 hours. Evidence of a gopher's burrow system includes mounds, soil plugs, and winter soil casts.

### Mounds

As a pocket gopher tunnels, it loosens soil with its front legs. When digging becomes difficult, it bites off chunks of earth or roots with its incisors. The gopher then somersaults to turn around and push the loose earth and other debris to the surface bulldozer style, with its front feet and head. The excavated material is pushed out of the exit tunnel to the front, right and left, creating a fan-shaped or heart-shaped mound (Fig. 3). (See Table 1 for a comparison of pocket gopher and mole mounds.)

The capacity of pocket gophers to excavate tunnels is phenomenal and heavy clay soil doesn't deter them. Gophers may create several mounds per day, especially during the seasons when the soil is moist and easy to dig. In irrigated areas such as lawns, gardens, and pastures, digging conditions may be optimal year-round, and mounds can appear at any time.

### Tunnels


Pocket gopher tunnels are  $1\frac{3}{4}$  to  $3\frac{1}{2}$  inches in diameter, depending on the size of the gopher digging the tunnel. Tunnels occur 4 to 12 inches below ground, whereas the nest and food storage chamber may be as deep as 6 feet. Tunnels tend to be deeper in drier soils. Short, sloping, lateral tunnels connect the main tunnel system to the surface and are created for pushing dirt to the surface and access to foraging on the surface (Fig. 3).

### Soil Plugs

Tunnel exits made by a pocket gopher are marked by a 1- to 3 inch circle of disturbed soil, or a circular depression, called a "soil plug." Soil plugs occur where a gopher emerged to forage or deposit soil, and then plugged up the hole on reentry. Plugs are found at mounds or along the course of the burrow system. Vegetation may be clipped around the soil plugs where a gopher was foraging.

### Winter Soil Casts

Soil casts are created because pocket gophers commonly backfill their previously excavated tunnels with excess soil when they dig new tunnels. Casts are the result of this excess soil being backfilled into snow tunnels. When



**Figure 3.** *The side view of a portion of a pocket gopher's burrow system showing the mound, the short, sloping lateral tunnel, and the deeper main tunnel.*

*(Drawing by Jenifer Rees.)*

Table 1. Differences between moles and pocket gophers.

Before setting out to control what you assume to be gopher damage, be sure to properly identify the animal causing the damage.

<b>Moles</b>	<b>Pocket Gophers</b>
Minute eyes are often not visible.	Small eyes are clearly visible.
Muzzle is long and tapering.	Muzzle is rounded.
The many small teeth are not apparent.	Orange, chisel-like pairs of upper and lower incisors are apparent.
Mounds are round when viewed from above.	Mounds are crescent- or heart-shaped when viewed from above.
Soil plug is in the middle of mound and may not be distinct.	Soil plug is in the middle of the V shape or off to the side of the mound and may leave a visible depression.
Tunnels are often just beneath the surface, leaving a raised ridge.	No tunnels are visible from above ground.

the snow melts, these then become apparent. Castings are nearly always fragmented and in short sections. Only a fraction of snow tunnels are backfilled with soil, so castings represent only a fraction of the gopher’s winter work.

## Preventing Conflicts

The ecological services of pocket gophers, which are substantial, are often not appreciated, particularly when the animals make their presence known by eating garden crops or damaging orchard or ornamental trees.

For homeowners and gardeners, gophers may be only an occasional (or seasonal) nuisance in lawns and garden beds, and not a long-term problem or threat. Where these animals are not so numerous as to be causing heavy damage, they should be considered neutral.

The subspecies **Brush Prairie pocket gopher** and the **Mazama (Western) pocket gopher** are in decline and are, or soon will be, of conservation concern (see “Legal Status” and “Mazama Pocket Gopher Conservation”). The presence of one of these species in an area where you plan to take action—chemical, nonchemical, mechanical, or otherwise—could preclude use of this action. Before moving forward with any type of control, contact your local Fish and Wildlife office.

The following are suggestions for reducing conflicts. In cases where these methods are not practical, contact your local County Extension Agent or local Fish and Wildlife office for further information.

**Frightening devices and repellents:** Although many devices are commercially available for use to frighten pocket gophers (vibrating stakes, ultrasonic devices, pinwheels, etc.), gophers do not frighten easily. This is probably because of their repeated exposure to noise and vibrations from sprinklers, people and livestock moving about, and lawnmowers and other power equipment. *Note:* Be skeptical of commercial products and claims, and make sure the manufacturer offers a money-back guarantee if the product proves ineffective.

No repellents currently available will reliably protect lawns or other plantings from pocket gophers. Mothballs, garlic, spearmint leaves, predator urine placed in tunnels—and a perimeter of mole plant or castor bean planted around gardens—have all provided mixed results. Such control strategies may be experimented with where gophers are an occasional problem, but not a long-term threat.

**Barriers:** Constructing a barrier to keep pocket gophers from tunneling into an area can be labor-intensive and costly; however, this approach is recommended for small areas and areas containing valuable plants. Flowerbeds and nursery beds can be protected by complete underground screening of the sides and bottom. Raised beds with rock or wooden side supports will only require bottom protection (Fig. 4).

Wire baskets can be used to protect the roots of individual trees and shrubs. These can be purchased from nurseries or farm supply centers, or be homemade. Use a double layer of light-gauge wire, such as 1-inch mesh chicken wire for trees and shrubs that will need protection only while young. Leave enough room to allow for a few years of root development before the wire rots away.

Groups of bulbs (gophers are reported not to eat daffodil bulbs) and other plants needing long-term protection can be placed in baskets made from 1/2-inch mesh hardware cloth, available from hardware stores and building supply centers.


Large areas, such as vegetable gardens, can be protected using an underground gopher fence (Fig. 5) or a stone-filled trench. However, such a below-ground barrier will only slow the movements of gophers for a time; sooner or later the barrier will be breached since gophers are capable of digging much deeper than 24 inches.

To add to the life of underground barriers, spray on two coats of rustproof paint before installation. Above-ground parts can also be painted to blend in. Always check for utility lines before digging in an area.

Several types of barriers (plastic tubes, one gallon plant containers) are effective at protecting aboveground parts of small plants, such as newly planted conifers.


Gophers may be deterred from chewing underground sprinkler lines or utility cables by surrounding them with 6 to 8 inches of coarse gravel 1 inch or more in diameter.

(In situations where gophers are gnawing on water lines or wires, or are burrowing into dams and dikes, refer to the handout on Muskrats for management recommendations.)



**Figure 4.** Raised beds can be protected from gopher damage by screening the bottom side with 1/2-inch mesh hardware cloth.

(Drawing by Jenifer Rees.)



**Figure 5.** A pocket gopher fence should be installed at least 24 inches below ground (or down to the hardpan or bedrock) and 6 inches above ground.

Use a 36-inch wide roll of 1/2-inch mesh hardware cloth.

Before placing the hardware cloth perpendicularly in the trench, the bottom 6 inches should be bent outward at a 90-degree angle.

Alternatively, fill a 10-inch wide by 24-inch deep trench with gravel that is 1 inch or more in diameter.

**Note:** Such barriers will likely only work temporarily.

(Drawing by Jenifer Rees.)

**Flooding:** Pocket gophers can easily withstand normal garden or home landscape irrigation, but flooding can sometimes be used to force them from their burrows. The entire tunnel system will need to be quickly and completely flooded to evict its tenets. Five-gallon buckets of water poured in the hole will flood the area more quickly than a running hose.

Flooding has the greatest chance of succeeding if gophers are invading the property for the first time. Where they are already well established, their systems are too extensive.

For humane reasons, concentrate this effort in late winter and early spring, before gophers give birth. Be careful when attempting to flood out a gopher near a building; doing so could damage the foundation or flood the basement or crawl space.

**Natural control:** A long-term way to help prevent conflicts is a combination of natural and active control. Predators—including snakes, dogs, coyotes, long-tailed weasels, and skunks—kill gophers. In addition, attracting barn owls and other raptors, which prey on young gophers when they disperse, may help control a gopher population, particularly in rural areas. Encouraging these species, or not discouraging them, may help control the gopher population.

Predators alone won't keep a gopher population below the levels that cause problems in gardens and landscaped areas. Before removing every gopher, they will move on to hunt at more profitable locations. However, when combined with the other control techniques described here, natural control can contribute to overall control.

## Public Health Concerns

Gophers are not considered to be a significant source of any infectious disease transmittable to humans or domestic animals.

## Legal Status

The subspecies **Brush Prairie pocket gopher** (*Thomomys talpoides douglasi*) of Clark County is a state candidate species, and the **Mazama (Western) pocket gopher** (*Thomomys mazama*) of Thurston, Pierce, and Mason Counties is a state threatened and federal candidate species. Because only remnant populations of these subspecies and species exist, people should not use lethal control in these areas.

## Mazama Pocket Gopher Conservation

In the south Puget Sound area, many populations of Mazama pocket gopher have disappeared since the 1940s, and the species was recently listed as a candidate for protection under the federal Endangered Species Act.

Mazama pocket gophers continue to decline in numbers in part because of their small, local breeding populations. For many years the species has persisted by continually recolonizing areas after local extinctions have occurred; however, loss of habitat to development, trapping by homeowners, and persecution by domestic cats and dogs have probably stopped much of this recolonization.

While large populations of some pocket gopher species can recover, the small and isolated populations of the Mazama pocket gopher can be completely lost.

If Mazama pocket gophers are to persist in the south Puget Sound area, they will require protection and lands where management is compatible with their needs. In addition, because Mazama gophers occupy grassy areas near homes and private property, a heightened level of tolerance will be required from those people who share their territories. In addition, if gophers are to survive in the suburbs, it may only be because homeowners are willing to keep their cats indoors.

The last records of Tacoma pocket gophers, *T.m. tacomensis* were of individuals killed by domestic cats (see the handout, "The Impacts of Domestic Dogs and Cats on Wildlife").

Elsewhere, pocket gophers are unclassified and may be trapped or killed and no special trapping permit is necessary for the use of live traps. However, a special trapping permit is required for the use of all traps other than live traps (RCW 77.15.192, 77.15.194; WAC 232-12-142). There are no exceptions for emergencies and no provisions for verbal approval. All special trapping permit applications must be in writing on a form available from the Department of Fish and Wildlife (WDFW).

It is unlawful to release a pocket gopher anywhere within the state, other than on the property where it was legally trapped, without a permit to do so (RCW 77.15.250; WAC 232-12-271).

Because legal status, trapping restrictions, and other information about gophers change, contact your local WDFW office for updates.

## Additional Information

### Books

Hygnstrom, Scott E., et al. *Prevention and Control of Wildlife Damage*. Lincoln, NE: University of Nebraska-Lincoln, Institute of Agriculture and Natural Resources, 1994. (Available from: University of Nebraska Cooperative Extension, 202 Natural Resources Hall, Lincoln, NE 68583-0819; phone: 402-472-2188; also see Internet Site below.)

Maser, Chris. *Mammals of the Pacific Northwest: From the Coast to the High Cascades*. Corvallis: Oregon State University Press, 1998.

Verts, B. J., and Leslie N. Carraway. *Land Mammals of Oregon*. Los Angeles: University of California Press, 1998.

### Internet Resources

Burke Museum's Mammals of Washington

Adapted from "Living with Wildlife in the Pacific Northwest" (see <http://wdfw.wa.gov/wlm/living.htm>)

Written by: Russell Link, Wildlife Biologist, [Linkrel@dfw.wa.gov](mailto:Linkrel@dfw.wa.gov)

Design and layout: Peggy Ushakoff, ITAS2

Illustrations: As credited

Copyright 2005 by the Washington Department of Fish and Wildlife.

This program receives Federal financial assistance from the U.S. Fish and Wildlife Service. It is the policy of the Washington State Department of Fish and Wildlife (WDFW) to adhere to the following: Title VI of the Civil Rights Act of 1964, Section 504 of the Rehabilitation Act of 1973, Title II of the Americans with Disabilities Act of 1990, the Age Discrimination Act of 1975, and Title IX of the Education Amendments of 1972. The U.S. Department of the Interior and its bureaus prohibit discrimination on the bases of race, color, national origin, age, disability and sex (in educational programs). If you believe that you have been discriminated against in any program, activity or facility, please contact the WDFW ADA Coordinator at 600 Capitol Way North, Olympia, Washington 98501-1091



